



## FACT SHEET

### Blast Lung Injury: What Clinicians Need to Know

Blast lung injury (BLI) presents unique triage, diagnostic, and management challenges and is a direct consequence of the blast wave from high explosive detonations upon the body. BLI is a major cause of morbidity and mortality for blast victims both at the scene and among initial survivors. The blast wave's impact upon the lung results in tearing, hemorrhage, contusion, and edema with resultant ventilation-perfusion mismatch. BLI is a clinical diagnosis and is characterized by respiratory difficulty and hypoxia, which may occur without obvious external injury to the chest.

Current patterns in worldwide terrorist activity have increased the potential for casualties related to explosions, yet few civilian health care providers in the United States have experience treating patients with explosion-related injuries. Emergency care providers are urged to learn more about the physics of explosions and other types of injuries that can result. Basic clinical information is provided here to inform practitioners of the presentation, evaluation, management, and outcomes of BLIs. Please see the reference list below for more information about how to treat injuries from explosions.

- **Clinical Presentation**

- Symptoms may include dyspnea, hemoptysis, cough, and chest pain.
- Signs may include tachypnea, hypoxia, cyanosis, apnea, wheezing, decreased breath sounds, and hemodynamic instability.
- Associated pathology may include bronchopleural fistula, air emboli, and hemothoraces or pneumothoraces.
- Other injuries may be present.

- **Diagnostic Evaluation**

- Chest radiography is necessary for anyone who is exposed to a blast. A characteristic "butterfly" pattern may be revealed upon x-ray.
- Arterial blood gases, computerized tomography, and doppler technology may be used.
- Most laboratory and diagnostic testing can be conducted per resuscitation protocols and further directed based upon the nature of the explosion (e.g. confined space, fire, prolonged entrapment or extrication, suspected chemical or biologic event, etc).



Photo courtesy of  
*Chest*, 1999.

- **Management**

- Initial triage, trauma resuscitation, treatment, and transfer should follow standard protocols; however some diagnostic or therapeutic options may be limited in a disaster or mass casualty situation.
- In general, managing BLI is similar to caring for pulmonary contusion, which requires judicious fluid use and administration ensuring tissue perfusion without volume overload.
- Clinical interventions
  - All patients with suspected or confirmed BLI should receive supplemental high flow oxygen sufficient to prevent hypoxemia (delivery may include non-rebreather masks, continuous positive airway pressure, or endotracheal intubation).

## Blast Lung Injury: What Clinicians Need to Know

(continued from previous page)

- Impending airway compromise, secondary edema, injury, or massive hemoptysis requires immediate intervention to secure the airway. Patients with massive hemoptysis or significant air leaks may benefit from selective bronchus intubation.
  - Clinical evidence of or suspicion for a hemothorax or pneumothorax warrants prompt decompression.
  - If ventilatory failure is imminent or occurs, patients should be intubated; however, caution should be used in the decision to intubate patients, as mechanical ventilation and positive end pressure may increase the risk of alveolar rupture and air embolism.
  - High flow oxygen should be administered if air embolism is suspected, and the patient should be placed in prone, semi-left lateral, or left lateral positions. Patients treated for air emboli should be transferred to a hyperbaric chamber.
- **Disposition and Outcome**
    - There are no definitive guidelines for observation, admission, or discharge following emergency department evaluation for patients with possible BLI following an explosion.
    - Patients diagnosed with BLI may require complex management and should be admitted to an intensive care unit. Patients with any complaints or findings suspicious for BLI should be observed in the hospital.
    - Discharge decisions will also depend upon associated injuries, and other issues related to the event, including the patient's current social situation.
    - In general, patients with normal chest radiographs and ABGs, who have no complaints that would suggest BLI, can be considered for discharge after 4-6 hours of observation.
    - Data on the short and long-term outcomes of patients with BLI is currently limited. However, in one study conducted on survivors one year post injury, no patients had pulmonary complaints, all had normal physical examinations and chest radiographs, and most had normal lung function tests.

Photo Source: Reprinted by permission from *Chest*. X-ray Figure I in "Recovery from Blast Lung Injury: One year follow-up", by Hirshberg, Boaz, MD, et al. Dec 1999, Vol 116(6), p 1683-88.

### References and Selected Readings:

- 1) Patterns of global terrorism 2001. United States Department of State. May 2002. United States Department of State website. Available at: <http://www.state.gov/documents/organization/10319.pdf>. Accessed 4 February 2005.
- 2) Patterns of global terrorism 2002. United States Department of State. April 2003. United States Department of State website. Available at: <http://www.state.gov/documents/organization/20177.pdf>. Accessed 4 February 2005.
- 3) Patterns of Global Terrorism 2003. United States Department of State. April 2004. United States Department of State website. Available at: <http://www.state.gov/documents/organization/31912.pdf>. Accessed 4 February 2005.
- 4) Terrorism 2000/2001. United States Department of Justice, Federal Bureau of Investigation, Counterterrorism Division. Publication #0308. Federal Bureau of Investigation website. Available at: [http://www.fbi.gov/publications/terror/terror2000\\_2001.htm](http://www.fbi.gov/publications/terror/terror2000_2001.htm). Accessed 4 February 2005.
- 5) Gadson LO, Michael ML, Walsh N (eds). FBI Bomb Data Center: 1998 Bombing Incidents, General Information Bulletin 98-1. US Department of Justice, Federal Bureau of Investigation. 1998.
- 6) Wightman JM, Gladish SL. Explosions and blast injuries. *Annals of Emergency Medicine*. 2001; 37(6): 664-678.
- 7) Horrocks CL Blast Injuries: Biophysics, pathophysiology, and management principles. *J R Army Med Corps* 2001;147:28-40.
- 8) Cullis, IG. Blast waves and how they interact with structures. *J R Army Med Corps* 2001;147:16-26.

July 7, 2005

Page 2 of 4

## Blast Lung Injury: What Clinicians Need to Know

(continued from previous page)

- 9) Langworthy MJ, Sabra J, Gould M. Terrorism and blast phenomena: lessons learned from the attack on the USS Cole (DDG67). *Clinic Orthop*. 2004; 422: 82-87.
- 10)Elsayed NM. Toxicology of Blast Overpressure. *Toxicology*. 1997; 121: 1-15.
- 11)Yetiser S and Ustun T. Concussive blast-type aural trauma, eardrum perforations, and their effects on hearing levels: an update on military experience in Izmir, Turkey. *Mil Med*. 1993; 158 (12):803-806.
- 12)Mayorga MA. The pathology of primary blast overpressure injury. *Toxicology*. 1997; 121:17-28.
- 13)Kluger Y. Bomb explosions in acts of terrorism-detonation, wound ballistics, triage, and medical concerns. *Isr Med Assoc J*. 2003; 5:235-240.
- 14)Sasser SM. Blast injuries. *The Turkish Journal of Emergency Medicine*. 2001; 1(1): 97-98.
- 15)Shaham, D, Sella T, Makori A, et al. The role of radiology in terror injuries. *Isr Med Assoc J*. 2002; 4(7): 564-567.
- 16)Knapp JF, Sharp RJ, Beatty R, et al. Blast trauma in a child. *Pediatric Emerg Care*. 1990; 6(2): 122-6.
- 17)Phillips YY. Primary blast injuries. *Ann Emerg Med*. 1986; 15:1446-1450.
- 18)Cooper GH, Maynard RL, Cross NL, et al. Casualties from terrorist bombings. *J Trauma*. 1983; 23(11): 955-967.
- 19)Stein M, Hirshberg A. Trauma care in the new millennium. *Surg Clin North Am*. 1999; 79 (6): 1537-1552.
- 20)Hadden WA, Rutherford WH, Merrett JD. The injuries of terrorist bombing: a study of 1532 consecutive patients. *Br J Surg*. 1978; 65(8):525-531.
- 21)Frykberg ER, Tepas JJ. Terrorist bombings: lessons learned from Belfast to Beirut. *Ann Surg*. 208: 569-576.
- 22)Mellor SG. The relationship of blast loading to death and injury from explosion. *World J Surg*. 1992; 16: 893-898.
- 23)Katz E, Ofek B, Adler J, et al. Primary blast injury after a bomb explosion on a civilian bus. *Ann Surg*. 1989; 209 (4): 484-488.
- 24)de Ceballos JPG, Turegano-Fuentes F, Perez-Diaz D, et al. 11 March 2004: The terrorist bomb explosions in Madrid, Spain-an analysis of the logistics, injuries sustained and clinical management of casualties treated at the closest hospital. *Crit Care Med*. 2005 Jan; 33(1):s107-112.
- 25)Pizov, R, Oppenheim-Eden A, Matot I, et al. Blast lung injury from an explosion on a civilian bus. *Chest*. 1999; 115 (1): 165-172
- 26)Tsokos M, Paulsen F, Petri S, Burkhard M, et al. Histological, immunohistochemical, and ultra-structural findings in human blast lung injury. *Am J Respir Crit Care Med*. 2003; 168: 549-555.
- 27)Leibovici D, Gofrit ON, Stein M, et al. Blast injuries: bus versus open-air bombings-a comparative study of injuries in survivors of open-air versus confined-space explosions. *J Trauma*. 1996; 41: 1030-1035.
- 28)Hirshberg B, Oppenheim-Eden A, Pizov R, et al. Recovery from Blast Lung Injury: One-Year Follow-up. *Chest*. 1999; 116(6): 1683-1688.
- 29)Frykberg ER, Tepas JJ, Alexander RH. The 1983 Beirut airport terrorist bombing: injury patterns and implications for disaster management. *Am Surg*. 1989; 55: 134-141.
- 30)Irwin RJ, Lerner MR, Bealer JF, Mantor PC, et al. Shock after blast wave injury is caused by a vagally mediated reflex. *J Trauma*. 1999; 47(1): 105-110.
- 31)Frykberg, ER. Medical management of disasters and mass casualties from terrorist bombings: how can we cope? *J Trauma*. 2002; 53:201-212.
- 32)Stuhmiller JH, Ho KHH, Vander Vorst MJ, et al. A model of blast overpressure injury to the lung. *J Biomech*. 1996; 29(2):227-234.
- 33)Stuhmiller JH. Biological response to blast overpressure: a summary of modeling. *Toxicology*. 1997; 121: 91-103.
- 34)Zhang J, Wang Z, Leng J, Yang Z. Studies on lung injuries caused by blast underpressure. *J Trauma*. 1996; 40 (3 supplement): s77-s80.

## Blast Lung Injury: What Clinicians Need to Know

(continued from previous page)

- 35) Elsayed NM, Gorbunov NV. Interplay between high energy impulse noise (blast) and antioxidants in the lung. *Toxicology*. 2003; 189(1-2):63-74.
- 36) Argyros, GJ. Management of Primary Blast Injury. *Toxicology*. 1997; 121: 105-115.
- 37) Leibovici D, Gofrit ON, Shapira SC. Eardrum perforation in explosion survivors: is it a marker of pulmonary blast injury?. *Ann Emerg Med*. 1999; 34(2): 168-172
- 38) Sorkine P, Szold O, Kluger Y, et al. Permissive hypercapnia ventilation in patients with severe pulmonary blast injury. *J Trauma*. 1998; 45(1): 35-38.
- 39) Irwin RJ, Lerner MR, Bealer JF, et al. Cardiopulmonary physiology of primary blast injury. *J Trauma*. 1997; 43 (4): 650-655.
- 40) Karmy-Jones R, Kissinger D, Golcovsky M. Bomb-related injuries. *Mil Med*. 1994; 159: 536-539.
- 41) Cohn SM. Pulmonary Contusion: review of the clinical entity. *J Trauma*. 1997 May; 42(5):973-79.
- 42) Elsayed NM, Gorbunov NV, Kagan VE. A proposed biochemical mechanism involving hemoglobin for blast over-pressure induced injury. *Toxicology*. 1997; 121: 81-90.
- 43) Lavonis E. Blast injuries. Emedicine website. Available at: <http://www.emedicine.com/emerg/topic63.htm>. Accessed 4 February 2005.
- 44) Gorbunov NV, McFaul SJ, Van Albert S, et al. Assessment of inflammatory response and sequestration of blood iron transferrin complexes in a rat model of lung injury resulting from exposure to low-frequency shock waves. *Crit Care Med*. 2004 Apr; 32(4): 1028-34.
- 45) DePalma RG, Burris DG, Champion HR, Hodgson MJ. Blast Injuries. *N Engl J Med* 2005; 352:1335-42.

For more information, visit [www.bt.cdc.gov](http://www.bt.cdc.gov),  
or call CDC at 800-CDC-INFO (English and Spanish) or 888-232-6348 (TTY).